

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 401/12, A61K 31/44, 31/55		A1	(11) International Publication Number: WO 99/24422
			(43) International Publication Date: 20 May 1999 (20.05.99)
(21) International Application Number: PCT/DK98/00478			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 5 November 1998 (05.11.98)			
(30) Priority Data: 1259/97 5 November 1997 (05.11.97) DK PA 1998 00850 26 June 1998 (26.06.98) DK			
(71) Applicant (for all designated States except US): NEUROSEARCH A/S [DK/DK]; Smedeland 26B, DK-2600 Glostrup (DK).			
(72) Inventors; and (75) Inventors/Applicants (for US only): PETERS, Dan [SE/SE]; Nygatan 17, S-232 31 Ärlöv (SE). OLSEN, Gunnar, M. [DK/DK]; (DK). NIELSEN, Simon, Feldbaek [DK/DK]; (DK). NIELSEN, Elsebet, Østergaard [DK/DK]; NeuroSearch a/s, Smedeland 26B, DK-2600 Glostrup (DK).			
(74) Common Representative: NEUROSEARCH A/S; Smedeland 26B, DK-2600 Glostrup (DK).			Published With international search report.

(54) Title: AZARING-ETHER DERIVATIVES AND THEIR USE AS NICOTINIC ACH RECEPTOR MODULATORS



(57) Abstract

The present invention discloses compounds represented by general formula (I) any of its enantiomers or any mixture thereof, or a pharmaceutically acceptable salt thereof; wherein m is 1, 2 or 3; n is 1, 2 or 3; R represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl or aralkyl; and R¹ represents a monocyclic 5 to 6 membered heterocyclic group which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, -COOR², -CONR²R³, -NH-CO₂R², NHCO-R², -OCO-NR²R³; wherein R² and R³ independently represents hydrogen or alkyl; aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; -X-alkyl-Y-alkyl wherein X and Y independently represents O, S, NH, N-alkyl or Se; and alkyl is optionally substituted with alkoxy or thioalkoxy; -X-(alkyl)_o-aryl- wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; -X-(alkyl)_o-5- or 6-membered heterocyclic monocyclic group wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; or R¹ represents a bicyclic heterocyclic group, composed of a 5 to 6 membered monocyclic heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, alkoxy-alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; and a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro. The compounds of the invention are useful as nicotinic ACh receptor ligands.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

AZARING-ETHER DERIVATIVES AND THEIR USE AS NICOTINIC ACh RECEPTOR MODULATORS

The present invention relates to novel pyridylether derivatives which are cholinergic ligands at nicotinic ACh receptors. The compounds of the invention are useful for the treatment of
5 condition or disorders or diseases involving the cholinergic system of the central nervous system, pain, inflammatory diseases, diseases caused by smooth muscle contractions and as assistance in the cessation of chemical substance abuse.

Background

10

The endogenous cholinergic neurotransmitter, acetylcholine, exert its biological effect via two types of cholinergic receptors; the muscarinic ACh receptors and the nicotinic ACh receptors. As it is well established that muscarinic ACh receptors dominate quantitatively over nicotinic ACh receptors in the brain area important to memory and cognition, much research aimed at
15 the development of agents for the treatment of memory related disorders have focused on the synthesis of muscarinic ACh receptor modulators. Recently, however, an interest in the development of nicotinic ACh receptor modulators has emerged. Several diseases are associated with degeneration of the cholinergic system i.e. senile dementia of the Alzheimer type, vascular dementia and cognitive impairment due to the organic brain damage disease
20 related directly to alcoholism. Indeed several CNS disorders can be attributed to a cholinergic deficiency, a dopaminergic deficiency, an adrenergic deficiency or a serotonergic deficiency. Alzheimer's disease is characterised by a profound loss of memory and cognitive functions caused by a severe depletion of cholinergic neurons, i.e. neurons that release acetylcholine. A reduction in the number of nicotinic ACh receptors are also observed with the progression of
25 Alzheimer's disease. It is believed that the neurons in the cortex that die with the progression of Alzheimer's disease do so because of lack of stimulation of the nicotinic ACh receptors. It is predicted that treatment of Alzheimer's patients with nicotinic ACh receptor modulators will not only improve the memory of patients but in addition act to keep these neurons alive. Smoking actually seems to protect individuals against neurodegeneration and compounds behaving on
30 these receptor may very likely have a generally neuroprotective effect.

However degeneration of the cholinergic system is not limited to individuals suffering from i.e. Alzheimers disease but is also seen in healthy aged adults and rats. Therefore it is suggested that the cholinergic system is involved and partly responsible for the memory disturbances seen in aged animals and humans. Nicotine receptor modulator may therefore be useful in the
35 treatment of Alzheimer's disease, memory loss, memory dysfunction, AIDS-dementia, senile dementia or neurodegenerative disorders.

Parkinson's disease appears to involve degeneration of dopaminergic neurons. One symptom of the disease has been observed to be loss of nicotinic receptors associated with the dopaminergic neurons and possibly interfering with the process of release of dopamine. As sustained nicotine administration increases the number of receptors present, administration of
5 nicotine receptor modulators may ameliorate the symptoms of Parkinson's disease. Other condition or disorders or disease ascribed to deficiencies in the dopaminergic system is: drug addiction, depression, obesity and narcolepsy.

Tourette's syndrome is a neuropsychiatric disorder involving a range of neurological and behavioral symptoms. It is believed that neurotransmitter dysfunction is involved though the
10 pathophysiology is still unknown and that nicotine will be beneficial in the treatment of the disease (Devor et. al. The Lancet, vol. 8670 p. 1046, 1989)

Schizophrenia is a severe psychiatric illness. Neuroleptic compounds have been used in the treatment of the disease, the effect of the compounds is believed to be interaction in the dopaminergic system. Nicotine is proposed to be effective in the treatment of schizophrenia
15 (ie. Adler et. al. Biol. Psychiatry, Vol. 32, p. 607-616, 1992.)

Nicotine has been reported to have an effect on neurotransmitter release in several systems. Release of acetylcholine and dopamine by neurons upon administration of nicotine has been reported (J. Neurochem. vol. 43, 1593-1598, 1984) and release of norepinephrine by Hall et. al. (Biochem. Pharmacol. vol. 21, 1829-1838, 1972) Release of serotonin by Hery et. al. (Arch.
20 Int. Pharmacodyn. Ther. vol. 296. p. 91-97, 1977). Release of glutamate by Toth et. al (Neurochem. Res. vol. 17, p. 265-271, 1992)

The serotonin system and dysfunction's of the serotonergic system is believed to be involved in diseases or conditions or disorders like: anxiety, depression, eating disorders, obsessive compulsive disorder, panic disorders, chemical substance abuse, alcoholism, pain, memory
25 deficits and anxiety, pseudodementia, Ganser's syndrome, migraine pain, bulimia, obesity, pre-menstrual syndrome or late luteal phase syndrome, tobacco abuse, post-traumatic syndrome, social phobia, chronic fatigue syndrome, premature ejaculation, erectile difficulty, anorexia nervosa, disorders of sleep, autism, mutism or trichotillomania.

Nicotine improves concentration and task performance. Therefore compounds exhibiting
30 nicotine receptor modulating properties will be likely to be useful compounds in the treatment of learning deficit, cognition deficit, attention deficit, attention deficit hyperactivity disorder and dyslexia.

Tobacco use and especially cigarette smoking is recognised as a serious health problem. However nicotine withdrawal symptoms associated with smoking cessation makes it difficult to
35 break this habit. Withdrawal symptoms include anger, anxiety, difficulties in concentrating,

restlessness, decreased heart rate and increased appetite and weight gain. Nicotine itself has shown to ease the withdrawal symptoms.

Withdrawal from addictive substances, i.e. opiates, benzodiazepines, ethanol, tobacco or
5 nicotine, is in general a traumatic experience characterised by anxiety and frustration. Nicotine has been found to be effective in reducing anger, irritability, frustration and feelings of tension without causing general response depression, drowsiness or sedation and compounds having same characteristics as nicotine is likely to have same effects.

10 Mild to moderate pain is normally treatable with NSAID's (non-steroidal anti-inflammatory drugs) while opiates are used preferentially for moderate to severe pain. The opiates have some well-known side-effects, including chemical dependence and abuse potential as well as a depressive effect on the respiratory and gastrointestinal system. There exists therefore a strong need for analgesic compounds that do not exhibit these side effects and which can
15 relieve mild, moderate and severe pain of acute, chronic or recurrent character as well as migraine pain and postoperative pain, phantom limb pain.

Epibatidine, a compound isolated from the skin of a poison frog, is a very potent analgesic with an approximate potency of 500 times that of morphine. The analgesic effect is not affected by naloxone, which is an indication of a negligible affinity for the opiate receptors. Epibatidine is
20 an nicotinic cholinergic receptor agonist and it is therefore very likely, that compounds possessing this receptor modulating character will also show a strong analgesic response. The compounds of the present invention has proven useful for modulation of smooth muscle contractions and may therefore be used in the treatment or prevention of condition or disorders or diseases inherent from smooth muscle contractions like i.e. convulsive disorders,
25 angina pectoris, premature labor, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia.

Further, it is well known that nicotine has an effect on appetite and it is predicted that modulators at the nicotine ACh receptor may be useful as appetite suppressants in the
30 treatment of obesity and eating disorders.

The cholinergic receptors play an important role in the functioning of muscles, organs and generally in the central or peripheral system. There are also complex interactions between cholinergic receptors and the function of receptors of other neurotransmitters such as dopamine, serotonin and noradrenaline.

35 It is likely that nicotine receptor modulator compounds can be effective in preventing or treating conditions or disorders or diseases like: inflammation, inflammatory skin conditions,

Chron's disease, inflammatory bowel disease, irritable colon, ulcerative colitis, irritable colon, diarrhoea, neurodegeneration, peripheral neuropathy, amyotrophic lateral sclerosis, nociception, endocrine disorders, thyrotoxicosis, pheochromocytoma, hypertension, arrhythmias, mania, manic depression, Huntington's disease, jetlag.

5

The compounds of the present invention are nicotine receptor modulators and has the potential to exhibit nicotinic pharmacology, preferentially without the side effects associated with nicotine itself. Additionally, the compounds are expected to have the potential as enhancers of neurotransmitter secretion and suppress symptoms associated with a low activity
10 of neurotransmitters.

Objects of the Invention

It is an object of the present invention to provide novel pyridylether derivatives which are
15 useful for the treatment of a range of diseases and disorders characterised by decreased cholinergic function or responsive to the activity of nicotinic ACh receptor agonists.

Another object of the present invention is to provide novel pharmaceutical compositions containing these compounds, as well as methods for the preparation thereof and methods for
20 the treatment therewith.


Other objects will become apparent hereinafter to one skilled in the art.

Summary of the Invention

25

The invention then, inter alia, comprises the following, alone or in combination:

A compound having the formula,



30

any of its enantiomers or any mixture thereof, or a pharmaceutically acceptable salt thereof;

wherein

m is 1,2 or 3;

5 n is 1, 2 or 3;

R represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl or aralkyl; and

R¹ represents

a monocyclic 5 to 6 membered heterocyclic group which may be substituted one or more
10 times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, -COOR³, -CONR²R³, -NH-CO₂R², NHCO-R², -OCO-NR²R³; wherein R² and R³ independently represents hydrogen or alkyl;

aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl
15 alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

X-alkyl-Y-alkyl wherein X and Y independently represents O, S, NH, N-alkyl or Se; and alkyl is optionally substituted with alkoxy or thioalkoxy;

-X-(alkyl)_o-aryl - wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se;
20 optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

-X-(alkyl)_o-5- or 6-membered heterocyclic monocyclic group wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃,
25 OCF₃, CN, amino and nitro;

- a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; or

30 or R¹ represents a bicyclic heterocyclic group, composed of a 5 to 6 membered monocyclic heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, alkoxy-alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl,
35 alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; and a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with

alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

The compounds as above may contain isotopes of one or more of the atoms present.

5

a pharmaceutical composition, comprising a therapeutically effective amount of a compound of above or a pharmaceutically acceptable addition salt thereof, together with at least one pharmaceutically acceptable carrier or diluent.

- 10 the use of a compound as above for the manufacture of a medicament for the treatment or prevention of a condition or disorder or disease of a living animal body, including a human, which condition or disorder or disease is responsive to the activity of nicotinic ACh receptor modulators;
- 15 a method of treating a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators, comprising the step of administering to such a living animal body, including a human, in need thereof a therapeutically effective amount of a compound as above;

20

Detailed disclosure of the invention

In a preferred embodiment of the invention the chemical compound of formula (I) an azetidine, pyrrolidine, piperidine or a homopiperidine is the group bridging R and R¹;

25

In another preferred embodiment of the invention

R is hydrogen or C₁₋₄-alkyl;

and R¹ represents a pyridyl or a substituted pyridyl;

- 30 In a more preferred embodiment,

R¹ is 3-pyridyl, 5-chloro-3-pyridyl, 2-chloro-3-pyridyl;

In another preferred embodiment the chemical compound of the invention is a compound of the general formula (I) said compound being:

- 35 (±)-1-Methyl-3-oxy-(3-pyridyl)-pyrrolidine;
(±)-1-Methyl-3-oxy-(3-pyridyl)-azetidine;

- (±)-1-Methyl-3-oxy-(3-pyridyl)-piperidine;
- (±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-pyrrolidine;
- (±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-piperidine;
- (±)-1-Methyl-3-oxy-(2-chloro-3-pyridyl)-piperidine;
- 5 (±)-1-Methyl-4-oxy-(3-pyridyl)-homopiperidine;
- (±)-1-Methyl-4-oxy-(5-chloro-3-pyridyl)-homopiperidine;
- (±)-3-Oxy-(3-pyridyl)-pyrrolidine;
- (±)-3-Oxy-(3-pyridyl)-azetidine
- (±)-3-Oxy-(3-pyridyl)-piperidine;
- 10 (±)-(5-Chloro-3-pyridyl)-3-oxy-pyrrolidine;
- (±)-(5-Chloro-3-pyridyl)-3-oxy-piperidine;
- (±)-(2-Chloro-3-pyridyl)-3-oxy-piperidine;
- (±)-4-Oxy-(3-pyridyl)-homopiperidine; or
- (±)-(5-Chloro-3-pyridyl)-4-oxy-homopiperidine;
- 15 any of its enantiomers or any mixture thereof, or a pharmaceutically acceptable salt thereof;

Pharmaceutical acceptable addition salts

The chemical compound of the invention may be provided in any form suitable for the intended
 20 administration. Suitable forms include pharmaceutically (i.e. physiologically) acceptable salts, and pre- or prodrug forms of the chemical compound of the invention.

Examples of pharmaceutically acceptable addition salts include, without limitation, the non-toxic inorganic and organic acid addition salts such as the acetate derived from acetic acid, the aconate derived from aconitic acid, the ascorbate derived from ascorbic acid, the
 25 benzenesulfonate derived from benzenesulfonic acid, the benzoate derived from benzoic acid, the cinnamate derived from cinnamic acid, the citrate derived from citric acid, the embonate derived from embonic acid, the enantate derived from enanthic acid, the formate derived from formic acid, the fumarate derived from fumaric acid, the glutamate derived from glutamic acid, the glycolate derived from glycolic acid, the hydrochloride derived from hydrochloric acid, the
 30 hydrobromide derived from hydrobromic acid, the lactate derived from lactic acid, the maleate derived from maleic acid, the malonate derived from malonic acid, the mandelate derived from mandelic acid, the methanesulfonate derived from methane sulphonic acid, the naphthalene-2-sulphonate derived from naphthalene-2-sulphonic acid, the nitrate derived from nitric acid, the perchlorate derived from perchloric acid, the phosphate derived from phosphoric acid, the
 35 phthalate derived from phthalic acid, the salicylate derived from salicylic acid, the sorbate derived from sorbic acid, the stearate derived from stearic acid, the succinate derived from

succinic acid, the sulphate derived from sulphuric acid, the tartrate derived from tartaric acid, the toluene-p-sulphonate derived from p-toluene sulfonic acid, and the like. Such salts may be formed by procedures well known and described in the art.

Other acids such as oxalic acid, which may not be considered pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining a chemical compound of the invention and its pharmaceutically acceptable acid addition salt.

Metal salts of a chemical compound of the invention includes alkali metal salts, such as the sodium salt of a chemical compound of the invention containing a carboxy group.

The chemical compound of the invention may be provided in solved or unsolved form together with a pharmaceutically acceptable solvents such as water, ethanol and the like. In general, solved forms are considered equivalent to dissolved forms for the purposes of this invention.

Steric Isomers

15 The chemical compounds of the present invention may exist in (+) and (-) forms as well as in racemic forms. The racemates of these isomers and the individual isomers themselves are within the scope of the present invention.

Racemic forms can be resolved into the optical antipodes by known methods and techniques.

One way of separating the diastereomeric salts is by use of an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optical active matrix. Racemic compounds of the present invention can thus be resolved into their optical antipodes, e.g., by fractional crystallisation of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example.

25 The chemical compounds of the present invention may also be resolved by the formation of diastereomeric amides by reaction of the chemical compounds of the present invention with an optically active activated carboxylic acid such as that derived from (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic acid or by the formation of diastereomeric carbamates by reaction of the chemical compound of the present invention with an optically active chloroformate or the like.

Additional methods for the resolving the optical isomers are known in the art. Such methods include those described by *Jaques J, Collet A, & Wilen S* in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Moreover, some of the chemical compounds of the invention contains double bonds and may thus exist in two forms, cis- and trans-forms (Z- and E-form), depending on the arrangement

of the substituents around the $-C=C-$ double bond. A chemical compound of the present invention may thus be the cis- or trans-form (Z- and E-form), or it may be a mixture hereof.

Definition of substituents

5

Halogen is fluorine, chlorine, bromine or iodine.

Alkyl means a straight chain or branched chain of one to eight carbon atoms, including but not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, and hexyl;

10

Cycloalkyl means cyclic alkyl of three to seven carbon atoms, including but not limited to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl;

Alkenyl means a group of from two to six carbon atoms, including at least one double bond, for example, but not limited to ethenyl, 1,2- or 2,3-propenyl, 1,2-, 2,3-, or 3,4-butenyl.

Alkynyl means a group of from two to six carbon atoms, including at least one triple bond, for example, but not limited to ethynyl, 1,2- or 2,3-propynyl, 1,2- or 2,3- or 3,4-butyne.

Cycloalkylalkyl means cycloalkyl as above and alkyl as above, meaning for example, cyclopropylmethyl.

Alkoxy is O-alkyl, wherein alkyl is as defined above.

Cycloalkoxy is O-cycloalkyl, wherein cycloalkyl is as defined above.

Alkenoxy is O-alkenyl wherein alkenyl is as defined above.

Alkynoxy is O-alkynyl, wherein alkynyl is as defined above;

30

Thioalkoxy is S-alkyl, wherein alkyl is as defined above.

Amino is NH_2 or NH -alkyl or N -(alkyl)₂, wherein alkyl is as defined above.

A monocyclic 5- to 6-membered heterocyclic group includes, for example, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl,

isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,5-oxadiazol-3-yl, 1,2,5-oxadiazol-4-yl, 1,2,5-thiadiazol-3-yl, 1,2,5-thiadiazol-4-yl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrazinyl and 3-pyrazinyl and 1-pyrazolyl, 3-pyrazolyl, and 4-pyrazolyl.

A bicyclic heterocyclic group composed of a 5 to 6 membered monocyclic heterocyclic group and a fused benzene ring means a monocyclic 5 to 6 membered heterocyclic group as above which is fused to a benzene ring including, for example, 2-, 3-, 4-, 5-, 6-, 7-benzofuranyl, 1-, 2-, 4-, 5- benzimidazolyl, 2-, 3-, 4-, 5-, 6-, 7-indolyl, 2-, 3-, 4-, 5-, 6-, 7-, 8-quinolinyl and 1-, 3-, 4-, 5-, 6-, 7-, 8-isoquinolinyl.

Aryl is an aromatic hydrocarbon, such as phenyl and naphthyl.

Aralkyl means alkyl as above and aryl as above, meaning for example benzyl, phenethyl.

Isotopes means one or more atom in the compound is substituted with an isotope of the naturally occurring atoms and includes though not limited to deuterium, tritium, ^{13}C , ^{14}C , ^{131}I , ^{125}I , ^{123}I , ^{18}F ;

Further, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

It will be appreciated by those skilled in the art that the compounds of the present invention may contain several chiral centers and that such compounds exist in the form of isomers (i.e. enantiomers). The invention includes all such isomers and any mixtures thereof including racemic mixtures.

Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can thus be resolved into their optical

antipodes, e.g., by fractional crystallization of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example. The compounds of the present invention may also be resolved by the formation of diastereomeric amides by reaction of the compounds of the present invention with an optically active activated carboxylic acid such as that derived from
5 (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic acid or by the formation of diastereomeric carbamates by reaction of the compounds of the present invention with an optically active chloroformate or the like.

Additional methods for the resolution of optical isomers, known to those skilled in the art may
10 be used, and will be apparent to the average worker skilled in the art. Such methods include those discussed by J. Jaques, A. Collet, and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Optical active compounds can also be prepared from optical active starting materials.
15

The compounds of the invention may be prepared by any conventional method useful for the preparation of analogous compounds and as described in the examples below.

Starting materials for the processes described in the present patent application are known or
20 can be prepared by known processes from commercially available materials

A compound of the invention can be converted to another compound of the invention using conventional methods.

25 The products of the reactions described herein are isolated by conventional means such as extraction, crystallization, distillation, chromatography, and the like.

Biology

30 Nicotinic ACh receptors in the brain are pentameric structures composed of subunits distinct from those found in skeletal muscles. The existence of seven α -subunits ($\alpha 2$ - $\alpha 7$, $\alpha 9$) and three β -subunits ($\beta 2$ - $\beta 4$) in the mammalian brain has been described.

The predominant subtype with high affinity for nicotine is comprised of $\alpha 4$ and $\beta 2$ subunits.
35

The affinity of compounds of the invention for nicotinic ACh receptors have been investigated in three test for *in vitro* inhibition of ^3H -epibatidin binding, ^3H - α -bungarotoxin binding and ^3H -cytisine binding as described below:

5 *In vitro* inhibition of ^3H -cytisine binding

The predominant subtype with high affinity for nicotine is comprised of α_4 and β_2 subunits. nAChRs of the latter type can selectively be labelled by the nicotine agonist ^3H -cytisine.

- 10 **Tissue Preparation:** Preparations are performed at 0-4°C unless otherwise indicated. Cerebral cortices from male Wistar rats (150-250 g) are homogenized for 20 sec in 15 ml Tris, HCl (50 mM, pH 7.4) containing 120 mM NaCl, 5 mM KCl, 1 mM MgCl_2 and 2.5 mM CaCl_2 using an Ultra-Turrax homogenizer. The homogenate is centrifuged at 27,000 x g for 10 min. The supernatant is discarded and the pellet is resuspended in fresh buffer and centrifuged a
15 second time. The final pellet is resuspended in fresh buffer (35 ml per g of original tissue) and used for binding assays.

- Assay:** Aliquots of 500 μl homogenate are added to 25 μl of test solution and 25 μl of ^3H -cytisine (1 nM, final concentration), mixed and incubated for 90 min at 2°C. Non-specific
20 binding is determined using (-)-nicotine (100 μM , final concentration). After incubation the samples are added 5 ml of ice-cold buffer and poured directly onto Whatman GF/C glass fibre filters under suction and immediately washed with 2 x 5 ml ice-cold buffer. The amount of radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

25

In vitro inhibition of ^3H - α -bungarotoxin binding Rat brain

- α -Bungarotoxin is a peptide isolated from the venom of the Elapidae snake *Bungarus multicinctus* (Mebs et al., Biochem. Biophys. Res. Commun., 44(3), 711 (1971)) and has high
30 affinity for neuronal and neuromuscular nicotinic receptors, where it acts as a potent antagonist. ^3H - α -Bungarotoxin binds to a single site in rat brain with an unique distribution pattern in rat brain (Clarke et al., J. Neurosci. 5, 1307-1315 (1985)).

- ^3H - α -Bungarotoxin labels nAChR formed by the α_7 subunit isoform found in brain and the α_1
35 isoform in the neuromuscular junction (Changeaux, Fidia Res. Found. Neurosci. Found. Lect. 4, 21-168 (1990). Functionally, the α_7 homo-oligomer expressed in oocytes has a calcium

permeability greater than neuromuscular receptors and, in some instances greater than NMDA channels (Seguela et al., J. Neurosci. 13, 596-604 (1993)).

Tissue preparation: Preparations are performed at 0-4°C unless otherwise indicated. Cerebral
5 cortices from male Wistar rats (150-250 g) are homogenized for 10 sec in 15 ml 20 mM Hepes
buffer containing 118 mM NaCl, 4.8 mM KCl, 1.2 mM MgSO₄ and 2.5 mM CaCl₂ (pH 7.5)
using an Ultra-Turrax homogenizer. The tissue suspension is centrifuged at 27,000 x g for 10
min. The supernatant is discarded and the pellet is washed twice by centrifugation at 27,000 x
g for 10 min in 20 ml fresh buffer, and the final pellet is resuspended in fresh buffer containing
10 0.01% BSA (35 ml per g of original tissue) and used for binding assays.

Assay: Aliquots of 500 µl homogenate are added to 25 µl of test solution and 25 µl of ³H- α -
bungarotoxin (2 nM, final concentration), mixed and incubated for 2 h at 37°C. Non-specific
binding is determined using (-)-nicotine (1 mM, final concentration). After incubation the
15 samples are added 5 ml of ice-cold Hepes buffer containing 0.05% PEI and poured directly
onto Whatman GF/C glass fibre filters (presoaked in 0.1% PEI for at least 6 h) under suction
and immediately washed with 2 x 5 ml ice-cold buffer. The amount of radioactivity on the filters
is determined by conventional liquid scintillation counting. Specific binding is total binding
minus non-specific binding.

20

***In vitro* inhibition of ³H-epibatidin binding**

Epibatidin is an alkaloid that was first isolated from the skin of the Ecuadoran frog
Epipedobates tricolor and was found to have very high affinity for neuronal nicotinic receptors,
25 where it acts as a potent agonist. ³H-epibatidin binds to two sites in rat brain, both of which
have pharmacological profiles consistent with neuronal nicotinic receptors and a similar brain
regional distribution (Hougling et al., Mol. Pharmacol. 48, 280-287 (1995)).

The high affinity binding site for ³H-epibatidin is most certainly binding to the $\alpha_4\beta_2$ subtype of
30 nicotinic receptors. The identity of the low affinity site is still unknown; does it represent a
second nicotinic receptor or a second site in the same receptor. The inability of α -bungarotoxin
to compete for ³H-epibatidin binding sites indicates that neither site measured represents the
nicotinic receptor composed of α_7 subunits.

35 Tissue preparation: Preparations are performed at 0-4°C unless otherwise indicated. The
forebrain (\pm cerebellum) from a male Wistar rat (150-250 g) is homogenized for 10- 20 sec in

20 ml Tris, HCl (50 mM, pH 7.4) using an Ultra-Turrax homogenizer. The tissue suspension is centrifuged at 27,000 x g for 10 min. The supernatant is discarded and the pellet is washed three times by centrifugation at 27,000 x g for 10 min in 20 ml fresh buffer, and the final pellet is resuspended in fresh buffer (400 ml per g of original tissue) and used for binding assays.

- 5 **Assay:** Aliquots of 2.0 ml homogenate are added to 0.100 ml of test solution and 0.100 ml of ^3H -epibatidin (0.3 nM, final concentration), mixed and incubated for 60 min at room temperature. Non-specific binding is determined using (-)-nicotine (30 μM , final concentration). After incubation the samples are poured directly onto Whatman GF/C glass fibre filters (presoaked in 0.1% PEI for at least 20 min) under suction and immediately washed with 2 x 5
- 10 ml ice-cold buffer. The amount of radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

Compound	^3H -	^3H -	^3H - α -
	cytisine $\text{IC}_{50}(\mu\text{M})$	epibatidin $\text{IC}_{50}(\mu\text{M})$	bungarotoxin $\text{IC}_{50}(\mu\text{M})$
1-Methyl-3-oxy-(3-pyridyl)-pyrrolidine (1a)	0,10	0,70	16,0
1-Methyl-3-oxy-(3-pyridyl)-piperidine(3a)	0,19	0,72	19,0
4-Oxy-(3-pyridyl)-homopiperidine fumaric acid salt (7b)	0,06	0,30	20,0
3-Oxy-(3-pyridyl)-pyrrolidine (1b)	0,07	0,41	>30

- The compounds of the present invention show high degree of binding affinity as well as
- 15 selectivity for one or more of the receptor subtypes. Therefore they have proven useful as selective receptor modulators of the nicotine receptor subtypes.

Pharmaceutical Compositions

- 20 While it is possible that, for use in therapy, a compound of the invention may be administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical formulation.

- The invention thus further provides pharmaceutical formulations comprising a compound of
- 25 the invention or a pharmaceutically acceptable salt or derivative thereof together with one or more pharmaceutically acceptable carriers therefor and, optionally, other therapeutic and/or prophylactic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Pharmaceutical formulations include those suitable for oral, rectal, nasal, topical (including buccal and sub-lingual), vaginal or parenteral (including intramuscular, sub-cutaneous and intravenous) administration or in a form suitable for administration by inhalation or insufflation.

5

The compounds of the invention, together with a conventional adjuvant, carrier, or diluent, may thus be placed into the form of pharmaceutical compositions and unit dosages thereof, and in such form may be employed as solids, such as tablets or filled capsules, or liquids such as solutions, suspensions, emulsions, elixirs, or capsules filled with the same, all for oral use, in
10 the form of suppositories for rectal administration; or in the form of sterile injectable solutions for parenteral (including subcutaneous) use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily
15 dosage range to be employed. Formulations containing ten (10) milligrams of active ingredient or, more broadly, 0.1 to one hundred (100) milligrams, per tablet, are accordingly suitable representative unit dosage forms.

The compounds of the present invention can be administrated in a wide variety of oral and
20 parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms may comprise, as the active component, either a compound of the invention or a pharmaceutically acceptable salt of a compound of the invention.

For preparing pharmaceutical compositions from the compounds of the present invention,
25 pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

30

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding
35 capacity in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid forms suitable for oral administration.

10

For preparing suppositories, a low melting wax, such as admixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

15

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

20 Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water-propylene glycol solutions. For example, parenteral injection liquid preparations can be formulated as solutions in aqueous polyethylene glycol solution.

The compounds according to the present invention may thus be formulated for parenteral administration (e.g. by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavours, stabilizing and thickening agents, as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, or other well known suspending agents.

- 5 Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavours, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

10

For topical administration to the epidermis the compounds according to the invention may be formulated as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents.

15

Formulations suitable for topical administration in the mouth include lozenges comprising active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

20

Solutions or suspensions are applied directly to the nasal cavity by conventional means, for example with a dropper, pipette or spray. The formulations may be provided in single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomising spray pump.

25

Administration to the respiratory tract may also be achieved by means of an aerosol formulation in which the active ingredient is provided in a pressurised pack with a suitable propellant such as a chlorofluorocarbon (CFC) for example dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by provision of a metered valve.

30
35

Alternatively the active ingredients may be provided in the form of a dry powder, for example a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP).

Conveniently the powder carrier will form a gel in the nasal cavity. The powder composition
5 may be presented in unit dose form for example in capsules or cartridges of, e.g., gelatin, or blister packs from which the powder may be administered by means of an inhaler.

In formulations intended for administration to the respiratory tract, including intranasal formulations, the compound will generally have a small particle size for example of the order of
10 5 microns or less. Such a particle size may be obtained by means known in the art, for example by micronization.

When desired, formulations adapted to give sustained release of the active ingredient may be employed.

15

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or
20 ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Tablets or capsules for oral administration and liquids for intravenous administration are preferred compositions.

25

Method of Treating

The compounds of the present invention are valuable nicotinic ACh receptor modulators and therefore useful for the treatment of a range of ailments involving cholinergic dysfunction as
30 well as a range of disorders responsive to the activity of nicotinic ACh receptor modulators.

The compounds may be used in the treatment, prevention, prophylaxis or alleviation of a disease, disorder or condition of the central or peripheral system as for example:
neurodegenerative disorders, cognitive or memory dysfunction, Alzheimer's disease,
Parkinson's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, Gilles de la
35 Tourettes syndrome, attention deficit hyperactivity disorder, anxiety, depression, mania, manic depression, schizophrenia, obsessive compulsive disorders, eating disorders like anorexia

nervosa, bulimia and obesity, narcolepsy, nociception, memory loss, memory dysfunction, AIDS-dementia, senile dementia, peripheral neuropathy, learning deficit, cognition deficit, attention deficit, autism, dyslexia, tardive dyskinesia, hyperkinesia, epilepsy, bulimia, post-traumatic syndrome, social phobia, chronic fatigue syndrome, disorders of sleep,
- 5 pseudodementia, Ganzer's syndrome, premenstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, premature ejaculation, erectile difficulty, mutism and trichotillomania.

The compounds of this invention may also be used in the treatment of inflammatory conditions
10 as for example: inflammatory skin conditions like acne and rosacea, Chron's disease, inflammatory bowel disease, ulcerative colitis, irritable colon, diarrhoea.

Also the compounds of the invention may be used in the treatment of diseases associated with smooth muscle contractions as for example: convulsive disorders, angina pectoris, premature
15 labor, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia.

The compounds of this invention may also be used in the treatment of pain as for example chronic, acute and recurrent pain, postoperative pain, migraine pain or phantom limb pain;
The compounds of the present invention may also be used for the assistance in cessation of
20 abuse of chemical substances as for example smoking cessation as well as cessation of use of other nicotine containing products, cessation of use of opioids like heroin, cocaine and morphine and cessation of use of benzodiazepines or alcohol. In the context of the present invention "treatment" means as well treatment as prevention, profylaxis and alleviation of
25 withdrawal symptoms and abstinence as well as treatment resulting in a voluntary diminished intake of the addictive substance.

Suitable dosage range are 0.1-500 milligrams daily, and especially 10-70 milligrams daily, administered once or twice a day, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the
30 subject involved and the body weight of the subject involved, and further the preference and experience of the physician or veterinarian in charge.

I.p. means intraperitoneally, which is a well known route of administration.

P.o. means peroral, which is a well known route of administration.

35

The invention then comprises the following alone or in combination:

the use of a compound as above wherein the disease to be treated is pain, a disease in the central nervous system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of
5 intake of the chemical substance.

The use as above wherein the disease is a disease in the central nervous system said disease being Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit
10 hyperactivity disorder.

The use as above wherein the disease to be treated is chemical substance abuse or withdrawal symptoms caused by the cessation of intake of the chemical substance, said chemical substance abuse being smoking or use of other nicotine containing products and withdrawal symptoms caused by cessation of use of nicotine containing products;
15

The method as above wherein pain, a disease in the central or peripheral system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of intake of chemical substances, such as smoking cessation, is treated.
20

The method as above wherein chemical substance abuse or withdrawal symptoms caused by the cessation of intake of the chemical substance, said chemical substance abuse being smoking or use of other nicotine containing products and withdrawal symptoms caused by cessation of use of nicotine containing products, is treated;
25

A method as above wherein a disease in the central or peripheral system, said disease being Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity disorder, is treated.

30 The following examples will illustrate the invention further, however, they are not to be construed as limiting.

Examples

General: All reactions involving air sensitive reagents or intermediates were performed under nitrogen and in anhydrous solvents. Magnesium sulfate was used as drying agent in the workup-procedures and solvents were evaporated under reduced pressure.

5 Method A

(±)-1-Methyl-3-oxy-(3-pyridyl)-pyrrolidine (1a)

A solution of 3-oxy-(3-pyridyl)-pyrrolidine (3.0 g, 18.3 mmol), formic acid (25.3 g, 99 %), formaldehyde (5.5 g, 37 %) and water (50 ml) was stirred at reflux for 15 h. The mixture was evaporated and sodium hydroxide (60 ml, 4 M) was added and the product was extracted two
10 times with ethyl acetate (60 ml). The product was isolated as an oil. Yield 2.5 g, 77 %.

(±)-1-Methyl-3-oxy-(3-pyridyl)-azetidine oxalic acid salt (2a)

Prepared according to method A. Mp. 166.9 - 170 °C.

15 (±)-1-Methyl-3-oxy-(3-pyridyl)-piperidine (3a)

Prepared according to method A. Isolated as an oil.

(±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-pyrrolidine (4a)

Prepared according to method A. Isolated as an oil.

20

(±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-piperidine fumaric acid salt (5a)

Prepared according to method A Mp 137.2 - 139.4 °C.

(±)-1-Methyl-3-oxy-(2-chloro-3-pyridyl)-piperidine (6a)

25 Prepared according to method A. Isolated as an oil.

(±)-1-Methyl-4-oxy-(3-pyridyl)-homopiperidine fumaric acid salt (7a)

Prepared according to method A. Mp 76 - 78 °C.

30 (±)-1-Methyl-4-oxy-(5-chloro-3-pyridyl)-homopiperidine fumaric acid salt (8a)

Prepared according to method A. Mp 113 - 116 °C.

Method B

35 (±)-3-Oxy-(3-pyridyl)-pyrrolidine (1b)

To a mixture of (\pm)-3-oxy-(3-pyridyl)-1-*tert*-butoxycarbonylpyrrolidine (16.0 g, 60.8 mmol) and dichloromethane (300 ml) was added trifluoroacetic acid (97 ml) and stirred for 6 h followed by evaporation. Methanol (50 ml) was added and the solution was cooled on ice. Sodium hydroxide (10 g, 0.25 mol) was added slowly and the mixture was stirred for 15 h. The solvent
5 volume was reduced to 10 ml and dichloromethane (30 ml) was added. The crude mixture was chromatographed on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as an oil. Yield 5.0 g, 50 %.

(\pm)-3-Oxy-(3-pyridyl)-azetidine trifluoroacetic acid salt (2b)

10 Prepared according to method B. Mp 151.8 - 152.7 °C.

(\pm)-3-Oxy-(3-pyridyl)-piperidine (3b)

Prepared according to method B. Isolated as an oil

15 **(\pm)-(5-Chloro-3-pyridyl)-3-oxy-pyrrolidine fumaric acid salt (4b)**

Prepared according to method B. Mp 141.7 - 143.2 °C.

(\pm)-(5-Chloro-3-pyridyl)-3-oxy-piperidine fumaric acid salt (5b)

Prepared according to method B. Mp 170.5 - 172.7 °C.

20

(\pm)-(2-Chloro-3-pyridyl)-3-oxy-piperidine fumaric acid salt (6b)

Prepared according to method B. Mp 143.9 - 146.0 °C.

(\pm)-4-Oxy-(3-pyridyl)-homopiperidine fumaric acid salt (7b)

25 Prepared according to method B. Mp 117 - 119 °C.

(\pm)-(5-Chloro-3-pyridyl)-4-oxy-homopiperidine (8b)

Prepared according to method B. Isolated as an oil.

30 **Method C**

(\pm)-3-Oxy-(3-pyridyl)-1-*tert*-butoxycarbonylpyrrolidine

Diethylazodicarboxylate (16.7 g, 96 mmol) was added dropwise to a mixture of tetrahydrofuran (150 ml) and triphenylphosphine (25.2 g, 96 mmol) and stirred for 0.5 h. (\pm)-3-Hydroxy-1-*tert*-butoxycarbonylpyrrolidine (12.0 g, 64 mmol) solved in tetrahydrofuran (50 ml) and added
35 dropwise, followed by 3-hydroxypyridine (9.1 g, 96 mmol). The mixture was stirred for 15 h at 40 °C. The solvent was evaporated and sodium hydroxide (200 ml, 1 M) was added followed

by extraction three times with diethyl ether (200 ml). The solvent was reduced to half volume by evaporation and triphenylphosphine oxide was filtered off. The crude mixture was chromatographed on silica gel with dichloromethane and methanol (25: 1). The title compound was obtained. Yield 16.1 g, 100 %.

5

(±)-4-Oxy-(3-pyridyl)-1-*tert*-butoxycarbonylhomopiperidine

Prepared according to method C.


1-*Tert*-butoxycarbonyl-azetidin-3-ol

- 10 A mixture of azetidin-3-ol (7.0 g, 96 mmol) (S. S. Chattarjee and D. J. Trigg Chem Commun. 93, (1968)), triethylamine (10.7 g, 106 mmol) and di-*tert*-butyl dicarbonat (21,0 g, 96 mmol) in dichloromethane (150 ml) was stirred for 15 h. The crude mixture was evaporated and chromatographed on silica gel with dichloromethane and ethanol (4 %). The title compound was obtained. Yield 12.9 g, 78 %.

15

Claims:

1. A compound represented by the general formula



5

any of its enantiomers or any mixture thereof, or a pharmaceutically acceptable salt thereof;

wherein

m is 1, 2 or 3;

10 n is 1, 2 or 3;

R represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl or aralkyl; and

R¹ represents

a monocyclic 5 to 6 membered heterocyclic group which may be substituted one or more
 15 times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, -COOR³, -CONR²R³, -NH-CO₂R², NHCO-R², -OCO-NR²R³; wherein R² and R³ independently represents hydrogen or alkyl;

aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl
 20 alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

-X-alkyl-Y-alkyl wherein X and Y independently represents O, S, NH, N-alkyl or Se; and alkyl is optionally substituted with alkoxy or thioalkoxy;

-X-(alkyl)_o-aryl - wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se;
 25 optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

-X-(alkyl)_o-5- or 6-membered heterocyclic monocyclic group wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;
 30

- a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; or
- 5 or R¹ represents a **bicyclic heterocyclic group**, composed of a 5 to 6 membered monocyclic heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, alkoxy-alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro; and a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;
- 15 2. A compound of formula (I), according to claim 1, wherein R and R¹ is attached to an azetidine, pyrrolidine, piperidine or homopiperidine;
3. A compound of formula (I), according to claim 1, wherein n is 1 and m is 1 or 2; R is hydrogen or C₁₋₄-alkyl;
- 20 and R¹ represents a pyridyl or a substituted pyridyl;
4. A compound of formula (I), according to claim 1, wherein R¹ is 3-pyridyl, 5-chloro-3-pyridyl, 2-chloro-3-pyridyl;
- 25 5. A compound of formula (I), according to claim 1, said compound being:
- (±)-1-Methyl-3-oxy-(3-pyridyl)-pyrrolidine;
- (±)-1-Methyl-3-oxy-(3-pyridyl)-azetidine;
- (±)-1-Methyl-3-oxy-(3-pyridyl)-piperidine;
- (±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-pyrrolidine;
- 30 (±)-1-Methyl-3-oxy-(5-chloro-3-pyridyl)-piperidine;
- (±)-1-Methyl-3-oxy-(2-chloro-3-pyridyl)-piperidine;
- (±)-1-Methyl-4-oxy-(3-pyridyl)-homopiperidine;
- (±)-1-Methyl-4-oxy-(5-chloro-3-pyridyl)-homopiperidine;
- (±)-3-Oxy-(3-pyridyl)-pyrrolidine;
- 35 (±)-3-Oxy-(3-pyridyl)-azetidine
- (±)-3-Oxy-(3-pyridyl)-piperidine;

- (±)-(5-Chloro-3-pyridyl)-3-oxy-pyrrolidine;
(±)-(5-Chloro-3-pyridyl)-3-oxy-piperidine;
(±)-(2-Chloro-3-pyridyl)-3-oxy-piperidine;
(±)-4-Oxy-(3-pyridyl)-homopiperidine; or
5 (±)-(5-Chloro-3-pyridyl)-4-oxy-homopiperidine;
or a pharmaceutically acceptable addition salt thereof;
6. A pharmaceutical composition, comprising a therapeutically effective amount of a compound of any of the claims 1 to 5, or a pharmaceutically acceptable addition salt thereof,
10 together with at least one pharmaceutically acceptable carrier or diluent.
7. The use of a compound according to any of the claims 1 to 5 for the manufacture of a medicament for the treatment of a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators.
15
8. The use according to claim 7 wherein the disease to be treated is pain, a disease in the central nervous system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of intake of the chemical substance such as smoking cessation;
20
9. The use of a compound according to claim 8 wherein a disease in the central nervous system is Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity disorder.
- 25 10. A method of treating a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators, comprising the step of administering to such a living animal body, including a human, in need thereof a therapeutically effective amount of a compound according to any of the claims 1 to 5.
- 30 11. The method according to claim 10, wherein pain, a disease in the central nervous system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of intake of the chemical substance, such as smoking cessation, are treated;
12. The method of claim 11 wherein a disease in the central nervous system, said disease
35 being Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity disorder, is treated.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 98/00478

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D401/12 A61K31/44 A61K31/55

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 28128 A (ZENECA LTD ;BROWN GEORGE ROBERT (GB); STOKES ELAINE SOPHIE ELIZABE) 7 August 1997 Compounds (IX) see claim 15	1
X	GB 2 295 387 A (GLAXO INC) 29 May 1996 R2 is (o)	1
X	DE 42 08 254 A (HOECHST AG) 16 September 1993 see claim 1; example 234	1
X	EP 0 351 282 A (SYNTHELABO) 17 January 1990 (III) in claim 2 see example 82	1
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 February 1999

Date of mailing of the international search report

26.02.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 98/00478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 338 331 A (BAYER AG) 25 October 1989 Compounds (II) see examples 32,33,84,85 ----	1
X	US 4 667 031 A (LILJE KENNETH C) 19 May 1987 See starting material of claim 1. see column 3, line 66 - line 67 ----	1
X	EP 0 149 088 A (DEGUSSA) 24 July 1985 Examples 1,2,4,7-10,22,49 see claim 1 ----	1
X	DE 32 27 741 A (HOECHST AG) 26 January 1984 see claim 1; examples 27-31; table 4 ----	1
X	BE 829 845 A (SOGERAS) 4 December 1975 see claim 1 ----	1
Y	US 5 629 325 A (LIN NAN-HORNG ET AL) 13 May 1997 see claim 1 ----	1-12
Y	WO 94 08992 A (ABBOTT LAB) 28 April 1994 see claim 1 ----	1-12
Y	WO 96 40682 A (ABBOTT LAB) 19 December 1996 see claim 1 -----	1-12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 98/00478

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 10-12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/ composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00478

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9728128	A	07-08-1997	AU 1608397 A EP 0880501 A	22-08-1997 02-12-1998
GB 2295387	A	29-05-1996	NONE	
DE 4208254	A	16-09-1993	AU 671108 B AU 3746693 A BR 9306083 A CA 2131545 A CN 1076692 A WO 9319050 A EP 0631575 A HU 67295 A JP 2619606 B JP 7506347 T KR 128270 B MX 9301399 A PL 175078 B US 5571815 A ZA 9301774 A	15-08-1996 21-10-1993 18-11-1997 30-09-1993 29-09-1993 30-09-1993 04-01-1995 28-03-1995 11-06-1997 13-07-1995 02-04-1998 01-11-1993 30-10-1998 05-11-1996 30-09-1993
EP 0351282	A	17-01-1990	FR 2634206 A FR 2640266 A AU 618378 B AU 3802989 A CA 1317292 A DE 68910210 D DE 68910210 T DK 341989 A ES 2062064 T FI 893365 A,B, IE 62216 B JP 2072173 A NO 174100 C PT 91126 A,B US 5034401 A	19-01-1990 15-06-1990 19-12-1991 18-01-1990 04-05-1993 02-12-1993 11-05-1994 25-04-1990 16-12-1994 13-01-1990 11-01-1995 12-03-1990 16-03-1994 08-02-1990 23-07-1991
EP 0338331	A	25-10-1989	DE 3835291 A AU 625817 B AU 3305989 A DK 186489 A ES 2045229 T GR 3006744 T IE 62064 B JP 1311059 A US 5453437 A US 5037841 A US 5274097 A	02-11-1989 16-07-1992 26-10-1989 20-10-1989 16-01-1994 30-06-1993 14-12-1994 15-12-1989 26-09-1995 06-08-1991 28-12-1993
US 4667031	A	19-05-1987	NONE	
EP 0149088	A	24-07-1985	AU 566560 B AU 3699684 A DD 231354 A DE 3443968 A DK 613384 A FI 845126 A,B GB 2152048 A,B	22-10-1987 04-07-1985 24-12-1985 31-10-1985 29-06-1985 29-06-1985 31-07-1985

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00478

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0149088 A		GR 82596 A	29-04-1985
		JP 60169476 A	02-09-1985
		PT 79736 B	30-12-1986
		SU 1417796 A	15-08-1988
		US 4643995 A	17-02-1987
DE 3227741 A	26-01-1984	NONE	
BE 829845 A	04-12-1975	FR 2306697 A	05-11-1976
		CA 1036597 A	15-08-1978
		CH 603626 A	31-08-1978
		DE 2528360 A	21-10-1976
		GB 1502182 A	22-02-1978
		JP 51118777 A	18-10-1976
		LU 72743 A	08-10-1975
		NL 7507441 A	12-10-1976
		US 3984411 A	05-10-1976
US 5629325 A	13-05-1997	WO 9746554 A	11-12-1997
WO 9408992 A	28-04-1994	AU 687017 B	19-02-1998
		AU 5355794 A	09-05-1994
		CA 2136075 A	28-04-1994
		EP 0663912 A	26-07-1995
		IL 107184 A	14-08-1997
		JP 8502481 T	19-03-1996
		MX 9306274 A	29-04-1994
WO 9640682 A	19-12-1996	AU 5804596 A	30-12-1996
		CA 2223062 A	19-12-1996
		EP 0846114 A	10-06-1998